

COMMUNITY Formulary by Clinical presentation UPDATED JUNE 2017 by the Tissue Viability Team.

See the Tissue Viability guidebook for advice on these presentations.

Limb Cellulitis

| | | |
|---|---|---|
| Dry Cellulitis | | Epimax or equivalent that is on the CCG emollients formulary |
| Wet cellulitis | Saturated enough to require daily dressings | Atrauman Zetuvit plus K-soft K- lite |
| Wet cellulitis | Saturated enough for 3 days dressing change | Adaptic touch Zetuvit E / or plus / or Kerramax Care k-soft K-lite |
| <p>Discussion: If open wound also then use antimicrobial Urgotul SSD as primary contact layer If so very wet also consider Kerramax Care as a primary dressing. To be used in conjunction with appropriate antibiotic regime. Compression should be used throughout treatment and pain assessment essential. Bandage toe to knee. If repeated cellulitis, patient to have full assessment to decrease risk.</p> | | |

Leaky Legs

Non-compression option

| | | |
|--|---|---|
| Legs saturated from below knees requiring intensive daily dressings management | Wash legs in Epimax If concerned for microbial load then consider Dermol Lotion for 2 weeks max (CCG formulary) | Zetuvit Plus or Kerramax Care as a primary dressing. K soft K-lite. |
|--|---|---|

Compression option

| | | |
|--|---|--|
| Legs saturated from below knees requiring intensive dressings management. Perform Doppler assessment | Wash legs in Epimax If concerned for microbial load then consider Dermol lotion 2 weeks max (CCG formulary) | Kerramax Care as a primary or if concerned for sticking add N-A ultra beneath. K soft to shape and protect, Actico short stretch bandage 8cm at foot then 10cm ankle to knee, 12cm knee to thigh (if required) Shaping necessary. |
| Toe bandaging. Use for leaking toes with chronic oedema, not if diabetic. | Apply following training only from Activa Healthcare or TVNS | Mollelast |

| | | |
|---|--|---|
| Microfine Toe caps. Caution with diabetes and ill-fitting causing pressure damage. Measure foot width to prescribe, | Ready to Wear washable compression toe caps (20-36mmHG) for toe and forefoot oedema. | XS,S,M,L,XL £26.92 www.hadhealth.com/forms |
| Cellona shoes (Activa Healthcare) to accommodate bandages | XS,S,,M,L,XL | On FP10 |
| Ensure there is a primary diagnosis e.g. dependency oedema, lymphoedema, venous eczema, cardiac failure and care plans will be dependent on this. | | |

Leg Ulcers

| | | |
|---|--|---|
| Venous leg ulcer with ABPI 0.8-1.3 See leg ulcer pathway | Superficial wounds – mix slough and granulation | Wash legs in Epimax If concerned for microbial load then consider Dermol Lotion for 2 weeks max (CCG formulary) N-A ultra or Adaptic Touch Kerramax Care to absorb, Actico if oedema Actico 2C if no oedema Or K-two range Prescribe according to ankle circumference |
| Venous leg ulcer with ABPI 0.8-1.3 | Sloughy wounds and deeper ulceration. TVN referral if complicated. | Gelling fibre to debride and absorb (Aquacel, Kytocel) Or Iodoflex if biofilm Kerramax Care to absorb and compression as above |
| Venous leg ulcers that are suitable for Hosiery kits (40mmHG) so smaller manageable leg ulcers | Measure and prescribe as per company info e.g. Activa leg ulcer kit, Actilymph leg ulcer kit Altipress (Urgo) Mediven ulcer kit | Consider a foam dressing under the hosiery kit (adhesive or non-adhesive) E,g Urgotul Absorb Border, Aquacel Foams |
| Mixed aetiology leg ulcers ABPI 0.5-0.8 (vascular referral) | Sloughy wounds and deeper ulceration | Aquacel to debride and absorb Kerramax Care and K-two reduced. |
| Arterial leg ulcers (vascular referral) | Usually sloughy, unlikely to debride despite optimal dressings | If painful then Urgotul or N/A Zetuvit Plus K-soft K-lite (no compression) If dry necrosis then Povitulle. |
| Complex compression Garments e.g. Farrowwrap, Readywraps . Microfine toe caps. | Washable compression wrap around devices. Expensive but a great option for self-care. | TVN referral Self-care /partnership care. |
| Healed leg ulcers | Educate patients on hosiery for maintenance for hosiery. Class 1 or 2 usually sufficient. | See below for all the companies |

| | | |
|-------|--|--------------------------|
| Shoes | | Cellona shoes if needed. |
|-------|--|--------------------------|

Venous eczema

| | | |
|-------------------|--|--|
| Wet Venous eczema | Compression will also resolve the eczema but wash legs and apply steroid creams and emollients to promote skin function. | Betamethasone Valerate 0.1% cream (until resolved approx. 2 weeks) applied daily, Epimax emollient, N-A Ultra, Kerramax Care and compression as per Doppler reading/ankle circumference. |
| Dry Eczema | Review compression status | Also consider Viscopaste but full assessment needed Betamethasone Valerate 0.1% cream or ointment for approx. 2 weeks until resolved. Encourage regular emollient application. |

Haematomas

| | | |
|------------------------------|---|---|
| Intact haematoma | Look at clotting status, refer to orthopaedic or plastics if suitable for surgical debridement. | Leave open and observe bruising is dispersing or for break in skin. May need emollient. |
| Broken and leaking haematoma | Check clotting status, refer to TVN, or orthopaedic or plastics if suitable for surgical debridement. Consider ABPI once debrided in view of compression. | Medihoney Antibacterial Wound gel to debride or Kerralite Cool Border or non-border to debride . Absorb and protect with Zetuvit E or Plus and K-soft , K-lite bandaging. |

Skin Tears

| | | |
|---|---|---|
| See pathway on guidebook. No skin loss or partial flap skin loss DO NOT apply any tape, adhesive strips, hydrocolloid, Iodine dressings, films or island dressing directly onto fragile skin. Use barrier films or spray (Derma S) under adhesive, if this cannot be avoided If venous disease present, treat | Low exudate levels Medium exudate levels | Urgotul Absorb Border Atrauman (2 days only , acute only) or Urgotul, Zetuvit E / K soft and K-lite Or Urgotul Absorb Border/ Mepilex Border |
|---|---|---|

| | | |
|----------------------|--|--|
| as venous leg ulcer. | | |
|----------------------|--|--|

Simple Wounds where healing is predicted.

| | | |
|---|------------------------------------|--|
| Low level exudate, uncomplicated wounds. May not suit very fragile thin skin. | 7 day wear time Showerproof | Cosmopore E dry dressing or Hydrofilm Plus (island dressing) |
|---|------------------------------------|--|

Surgical wounds

| | | |
|---|---|---|
| Healing by primary intention | Stitched or stapled | Hydrofilm Plus |
| Dehisced Rule out infection, consider surgical review. Must involve TVN. | Superficial Deep wound / cavity Infected Cavity | Urgotul Absorb Border / Mepilex Border Aquacel ribbon with Urgotul Absorb Border or TNP. Cutimed sorbact Ribbon |

Pressure Ulcers

| | | |
|------------------|------------------------------|---|
| Category 1 | No dressings needed, observe | Consider the use of Kerrapro over bony prominences. |
| Category 2 | Superficial | Hydrocoll or Urgotul Absorb Border |
| Category 3 and 4 | Superficial Cavity | Hydrocoll or Urgotul Absorb Border Aquacel Ribbon (AG if infected) or Kytocel, or Flaminal Hydro, Topical Negative Pressure (TVN referral) Urgotul Absorb Border sacrum or Granuflex Border sacrum |
| Unstageable | For active debridement | Medihoney antimicrobial gel or Kerralite Cool Border / non or Flaminal Hydro / Forte Larvae therapy Hydrocoll |
| Black Heels | If dry If broken skin | No dressing needed, measure, observe and elevate only. Do not apply foam dressings. As per podiatry recommends. Use Povitulle and Biatain non-adhesive, K-soft, K-lite. Refer for soft cast also. |

Moisture Lesions

| | | |
|--|---|---|
| Occasional urinary incontinence and faecal incontinence managed with regular pad changes | Intact, vulnerable with or without blanching redness | Ensure shaped pads are in use with adequate care package. Cleanse each pad change with PH balanced cleanser e.g. Epimax onto a cleansing cloth. This could also be the patient's own (PH balanced preferable). Apply derma S barrier cream if red skin from moisture. |
| Urinary and faecal incontinence with formed stool requiring regular pad changes | Blanching redness with sporadic rash maybe broken or not | Cleanse with Epimax or other emollient (from formulary) Derma- S barrier film applicators or spray, every three pad changes/ cleanses or daily. |
| Urinary and faecal incontinence with liquid stool requiring frequent pad changes | Blanching redness with sporadic rash and broken skin/ lesions | Derma- Pro Foam and spray incontinence cleanser. Apply Derma Pro skin protectant ointment onto the wound or broken skin (this is higher protection than the films)Do not use dressings. |

Fungating Wounds

| | | |
|---|---|--|
| Refer to TVN for individual prescribing ideas. For malodour – kills the anaerobes | This is a prescription only medicine, see BNF for prescribing details and contraindicators. | Anabact gel (Metronidazole 0.75%) |
| Other option gel for killing the anaerobes | | Flaminal forte gel for moderate to heavily exuding levels Flaminal Hydro – for low to moderate exuding levels |
| Charcoal dressings have mixed effectiveness | Charcoal options | Clinisorb |
| Primary dressings | Non-adherent wound contact layer | Adaptic Touch Urgotul |
| Alginates for haemostatic use | Be mindful these may stick and maybe more harmful to remove. | Kytocel |
| Secondary dressings | Choose secondary dressing as to anatomy and gentle removal. | Secure with Zetuvit Plus or Urgotul Absorb Border. Consider barrier creams and Comfifast if appropriate. |

Diabetic Foot or foot with a wound– Wound care to be led by PODIATRY

| | | |
|---|---|--|
| Wet diabetic ulcer cleansing | To Irrigate wet wounds only. Soak gauze in solution for at least 15 minutes | Prontosan wound irrigation solution |
| Wet or dry diabetic foot | Povitulle iodine (Inadine alternative) Conformable paste for infected or critically colonised wounds. | Povitulle Askina Caglitrol paste |
| Wet diabetic foot ulcers | Non adhesive foam due to maceration risk | Biatain non- adhesive |
| Wet diabetic foot ulcers | Alternative non- adhesive foam option as directed by podiatry. Can be cut to shape. | Mepilex XT |
| Black unstageable heel pressure ulcers that are dry | Following vascular assessment and cause of pressure damage | No dressing required |
| To secure foot dressings | | Hypafix Comfast red line Hospilite bandage |

Over granulating Wounds

Over granulation in wound beds is not always problematic, if wound measurements decrease then no action needed. Quite often moisture related.
 1st line – increase absorbency of dressing or increase frequency of dressing change
 If friable, bleeding and not responding then use an antimicrobial dressing for 2 weeks.
 Consider Hydrocortisone 1% cream to the over granulation as this may reduce inflammation in the tissue.
 Rule out BCC or Bowens disease also.

WOUND CARE FORMULARY 2017

Please inform the Tissue Viability Team if you wish to add any products to the formulary or want to give feedback regarding products on Tissue.referrals@nhs.net The formulary review group including CCG require for a justification form to be completed.

| CLEANSING AND EMOLLIENTS AND BARRIER PREPARATIONS | | |
|--|--|---|
| Tap water 1 st Line. | Potable tap water is safe to use for all wound care and surround skin cleansing but risk assess if infection risk and encourage clean drying. | Clinell clean and line bucket with bin liner if washing legs. |
| Irripod | 25 x 20mls | £5.84 |
| Prontosan Solution Ampules – as per podiatry instructions. Cleanser, removes biofilm and help to control bacterial levels on wounds | Soak gauze and apply to wound for 10 mins. | £14.18 for 24 |
| Octenisan wash Mitts – use as a leg wash alternative to water and whole body washing if needed (including hair) | Box of 10 1 pack per patient. Secure packet in between use. | 35p each |
| Epimax (see emollients formulary for product choices) | To use as a soap substitute /emollient 500g | £2.49 |
| Aquamax | To use as a soap substitute/emollient 500g | £3.99 |
| Zerocream | To use as a soap substitute/emollient 500g | £4.08 |
| Derma-Pro foam and spray incontinence cleanser | 250ml spray | £5.95 |
| Derma S Barrier Cream | 2g sachets (20) | £5.85 |
| | 28g | £2.98 |
| | 90g | £5.95 |
| Derma S Medical Barrier Film non-sting | Pump spray 30mls | £4.95 |
| | Foam Applicator 1ml (5) | £3.70 for 5 |
| Derma -Pro Skin Protectant ointment for Moisture lesions | 115g | £8.50 |
| DRESSING PACKS , TAPES AND TUBULAR BANDAGES | | |
| Nurse IT Small / medium Medium/large | For FP10 prescribing 7 x swabs, compartment tray, disposable forceps, 2 x laminated sterile fields, large apron, latex free gloves, measuring tape, towel and disposable bag. | 55p |
| Aseptic Nursing Pack medium Pack size 84 | EVX526 small EVX527 medium | 39p |

| | | |
|---|--|--|
| | EVX528 large | |
| Scanpor – skin friendly adhesive tape, For taping down dressing e.g. bandages and for skin contact. Multi- use tape | 2.5cm x 5m 5cm x 5m 7.5cm x 10m | 68p £1.18 £2.56 |
| Hypafix | 5cm x 5m 10cm x 5m | £1.45 £2.43 |
| Comfast Elasticated Viscose stockinette | 3.5cm x 1m (digits – red line) 7.5cm x 1m – blue line 7.5cm x 5m 10.75cm x 1m- yellow line 10.75cm x 5m 17.5 x 1m – beige line | 56p 77p £3.74 £1.20 £6.04 £1.83 |
| WOUND CONTACT LAYERS | | |
| Atrauman A non- adherent polyester mesh dressing, change every 3 days | 5 x 5cm 7.5 x 10cm 10 x 20cm 20 x 30cm | 27p 28p 63p £1.74 |
| Adaptic Touch A soft tack silicone mesh, 7 days wear time. Leave in situ and change secondary dressing more frequently. | 5 x 7.6cm 7.6 x 11cm 12.7 x 15cm 20 x 32cm | £1.13 £2.25 £4.65 £12.50 |
| N-A Ultra Knitted Viscose rayon sheet with a silicone coating 7 days wear time. | 9.5 x 9.5cm 9.5 x 19cm | 33p 63p |
| Urgotul A non- adherent ,flexible, contains TLC healing matric. 7 days wear time. Increases fibroblasts. | 5 x 5cm 10 x 10cm 15 x 15cm 15 x 20cm | £1.55 £3.11 £6.60 £8.79 |
| SUPER ABSORBERS | | |
| Zetuvit E Cellulose pad with fluid repellent packing. Change when saturated. Zetuvit E non- sterile e.g. for non- sterile areas such as groin, pilonidal sinuses | 10 x 10cm 10 x 20cm 20 x 20cm 20 x 40cm 10 x 10cm 10 x 20cm 20 x 20cm 20 x 40 cm | 21p 25p 39p £1.11 7p 9p 14p 27p |
| Zetuvit Plus Super absorbent fluid retaining particles, water repellent, soft Consistency. Change when saturated. | 10 x 10cm 10 x 20cm 15 x 20cm 20 x 25cm 20 x 40cm | 64p 88p £1.01 £1.38 £2.13 |

| | | |
|---|------------------------|-------|
| Kerramax Care Superabsorbent which locks in exudate. Horizontal wicking layer. For heavily exuding wounds. Suitable under compression. Change when saturated | 5 x 5cm | £1.00 |
| | 10 x 10cm | £1.29 |
| | 10 x 22cm | £1.68 |
| | 13.5 x 15.5 | £1.54 |
| | 20 x 22cm | £2.98 |
| | 20 x 30cm | £3.40 |
| | 20 x 50cm | £4.50 |
| | 21 x 23cm (multisite) | £3.23 |
| ISLAND DRESSINGS and FILMS – low absorbcy | | |
| Cosmopor E Self-adhesive island dressing with non- adherent absorbent pad. Not waterproof | 5 x 7.2cm | 8p |
| | 8 x 10cm | 18p |
| | 8 x 15cm | 28p |
| | 10 x 20cm | 46p |
| | 5 x 7.2cm | 18p |
| | 9x 10cm | 27p |
| Hydrofilm Plus Vapour permeable Adhesive film dressing with absorbent pad. Waterproof. | 9 x 15cm | 30p |
| | 10 x 20cm | 45p |
| | 10 x 30cm | 69p |
| | 6 x 7cm | 23p |
| | 10 x 12.5cm | 42p |
| | 10 x 15cm | 53p |
| Hydrofilm Vapour permeable adhesive film dressing with a high MVTR(Moisture Vapour Transmission Rate).Do not use in bleeding, infected or high exuding wounds. | 6 x 7cm | 23p |
| | 10 x 12.5cm | 42p |
| | 10 x 15cm | 53p |
| COMPRESSION BANDAGING - First line but other compression systems may be needed. Full vascular assessment essential prior to prescribing. | | |
| Actico 40+mmHg (variable) Inelastic compression bandage. To be used with full training on contraindications. Different application with ankle circumference above 25cm after wadding. | 8cm x 6m | £3.25 |
| | 10cm x 6m | £3.38 |
| | 12cm x 6m | £4.31 |
| Actico 2C Latex Free ulcer kit. Foam padding comfort layer and inelastic cohesive compression layer | Ankle 18-25cm | £8.03 |
| | Ankle25-32cm | £9.04 |
| Urgo K-Two (40mmHG) 2 layer bandage system of elastic and non-elastic components for graduated compression. Latex present | Ankle 18-25cm 10cm | £8.68 |
| | Ankle 25-32cm 10cm | £9.48 |

| | | |
|--|---|--|
| unless state latex free. Full training required for contraindications with compression. Prescribe by ankle circumference. | | |
| UrgoK-Two reduced (20mmHG) As above – order latex free | Ankle 18-25cm 10cm Ankle 25-32cm 10cm | £8.68 £9.48 |
| Mollelast (Activa Healthcare) Conforming bandage designed for dressing retention and finger and toe bandaging. Training required for toe bandage technique. | 4cm x 4m | 30p |
| Non- compression bandaging | | |
| K-soft- Absorbent non-woven sub bandage wadding. Viscose polyester, latex free. | 10cm x 3.5m 10cm x 4.5m | 46p 57p |
| K-Lite Lightweight knitted bandage type 2. Latex free. | 10cm x 4.5m 10cm x 5.25m | £1.01 £1.16 |
| Hospilite Cheaper alternative to K-lite type 2 light support bandage. | 7.5cm x 4.5m 10cm x 4.5m | 51p 62p |
| Viscopaste Zinc paste bandage for the management of chronic eczema, dermatitis under bandaging. Apply in pleats to avoid tourniquet. Address cause of eczema first. | 7.5cm x 6m | £3.69 |
| COMPRESSION HOSIERY – too many to put down individually so please contact the companies. To send product information. Measurement charts and prescribing options all on websites. | | |
| Activa | Actilymph hosiery kit Leg Ulcer Hosiery kit (40mmHG) Liner packs (10mmHG) | www.activahealthcare.co.uk also download app 08450606707 |
| Jobst | Ulcercare compression liner and stockings available | www.healthcare.co.uk/jobst-ulcercare-range 0207 7202266 |
| Urgo | Altipress 40 Leg ulcer Hosiery Kit Altipress liner packs Altimed Hosiery | www.altimed.co.uk 01509 501720 |
| Sigvaris | Ulcer X and other hosiery | www.sigvaris.com/uk 01264 326666 |
| Microfine Toe Caps (Haddenham healthcare) | Washable reusable compression garment for toes and forefoot. | £26.92 www.hadhealth.com/forms for measurements etc. |
| Medi 2 in 1 Donning / doffing Aide | Aids application of compression hosiery and armsleeves | £11.86 |
| Sockaid | Metal frame compression | £13.18 |

| | | |
|---|--|--|
| | garment applicator | |
| ANTIMICROBIAL DRESSINGS. | | |
| Aquacel Ag+ Extra Silver impregnated hydrofibre. Kills biofilm and prevents reformation and kills bacteria. 7 days wear time and review after 2 -4 weeks use. | 5 x 5cm 10 x 10cm 15 x 15cm 20 x 30cm 4 x 10cm | £1.95 £4.64 £8.75 £21.71 £2.83 |
| Povitulle (Inadine alternative) Non-adherent dressing containing 10% povidone iodine. Do not use if pregnant, breastfeeding, treating for kidney problems, before and after radio iodine, and patients with thyroid disease, burns or large injuries. | 5 x 5cm 9.5 x 9.5cm | 28p 42p |
| Iodoflex Cadexomer dressing with Iodine. For the treatment of small chronic exuding wounds and biofilms. Do not use on necrotic tissue. Do not use on children, pregnant or lactating women or people with thyroid or renal impairment. | 5g 10g | £4.13 £8.25 |
| Medihoney Antibacterial Wound gel. Medical grade honey with antibacterial and anti- inflammatory action. Use for superficial wounds, infected wounds, pressure ulcers, leg ulcers, malodourous, necrotic, surgical wounds. Do not use if sensitive to honey. Can be used for long term treatment. | 10g 20g | £2.69 £4.02 |
| Actilite Honey dressing Non-adherent viscose net dressing coated with 99% Manuka honey and 1% Manuka oil. Broad spectrum antimicrobial dressing. Can be used long term. Possible stinging sensation. | 5 x 5cm 10 x 10cm 10 x 20cm | 57p 98p £1.90 |
| KytoCel – Absorbent gelling fibre made from Chitosan, a | 5 x 5cm 10 x 10cm | 80p £1.92 |

| | | |
|---|--|---|
| broad spectrum antimicrobial. Debrides, removes in 1 x piece, Haemostatic properties. Aid autolytic debridement. For moderate to Highly exuding wounds. 7 days wear time. (replaces Sorbsan/ sorbsan silver) | 15 x 15cm 2.5 x 45cm | £3.60 £1.94 |
| Cutimed Sorbact Ribbon DACC coated antimicrobial ribbon designed to bind bacteria and fungi under moist wound conditions. For cavity or infected wounds. Suitable for fungal infection in the groin, skin folds or between digits. Cutimed sorbact swab As above but can be used folded or unfolded. 7 days wear but can be used for long term especially for re-infecting wounds. | 2 x 50cm 4x6cm (unfolded 11 x 16cm) 7 x 9cm(unfolded 17 x 27cm) | £4.03 £1.65 £2.75 |
| Askina Calgitrol Paste Conformable paste, ionic silver alginate for infected or critically colonised wounds. | 15g | £6.55 |
| Urgotul SSD A non-adherent antibacterial contact layer containing silver sulphadiazine particles. | 10 x 12cm 15 x 20cm | £3.17 £8.99 |
| ODOUR CONTROL | | |
| Anabact gel (POM) Metronidazole 0.75% gel for the deodorisation or malodorous Fungating tumours. | 15g | £4.47 |
| Clinisorb | 10 x 10cm 10 x 20cm 15 x 25cm | £1.91 £2.54 £4.09 |
| FOAMS | | |
| Urgotul Absorb Border Absorbent Foam dressing with silicone adhesive border.TLC wound contact layer, showerproof.7 days wear time | 8 x 8cm (incl Border) 8.8x15cm 10 x 10cm 10 x 25cm 13 x 13cm 15 x 20cm 20 x 20cm sacral | £1.36 £2.40 £1.96 £3.60 £2.40 £3.91 £4.27 |
| Mepilex Border Soft silicone flexible absorbent pad with film backing for moderate to highly exuding | 7x 7.5cm 10 x 10.5cm 10 x 12.5cm 15 x 17.5cm | £1.35 £1.90 £2.72 £4.18 |

| | | |
|---|--|---|
| wounds.7 days wear time | | |
| Mepilex XT Conformable soft silicone foam dressing for low and high viscosity. 7 days wear time. | 10 x 11cm 15 x 16cm | £2.66 £4.82 |
| Tegaderm Foam Adhesive Absorbent polyurethane foam dressing , 4 layers, oval shapes and spider dressing removal, good for contours e.g. heels. | 6.9x9.6cm (miniwrap) 6.9x7.6 (mini oval) 10 x 11cm (oval) 14.3 x 15.6 (oval) | £1.71 £1.46 £2.39 £4.24 |
| Biatain Non-Adhesive Soft conformable polyurethane foam dressing with vapour permeable film backing. Bevelled edges can be used under compression.1 st line for podiatry. | 5 x 7cm 10 x 20cm 10 x 10cm 15 x 15cm | £1.32 £3.96 £2.40 £4.43 |
| GELLING FIBRES AND PRODUCTS TO DEBRIDE | | |
| Aquacel Extra Hydrofibre which absorbs exudate and tansforms into a soft gel. Useful for debriding and absorbing, for all wet wounds. | 5 x 5cm 10 x 10cm 15 x 15cm 4 x 10cm | £1.00 £2.38 £4.48 £1.31 |
| Aquacel ribbon For cavity wounds. Stitched so removes in 1 piece. Document number of pieces used in a cavity. | 1 x 45cm ribbon 2 x 45cm ribbon | £1.86 £2.48 |
| Kytoceal Absorbent gelling fibre made from Chitosan, a broad spectrum antimicrobial (chitosan)Debrides, removes in 1 x piece, Haemostatic properites. Aids autolytic debridment. For moderate to highly exuding wounds. 7 days wear time. (replaces Sorbsan and sorbsan silver) | 5 x 5cm 10 x 10cm 15 x15cm 4 x 10cm 2.5 x 45cm ribbon | 80p £1.92 £3.60 £1.03 £1.94 |
| Kerralite Cool non- border Pro-ionic hydrogel contact layer and a fluid repellent polyurethane film outer layer to assist autolytic debridement of necrotic tissue and slough. For painful wounds, radiotherapy burns, burns and scalds. Not for full thickness, heavily bleeding wounds, 3 rd degree burns or as a covering | 6 x 6cm 12 x 8.5cm 18 x 12.5cm | £1.75 £2.57 £3.71 |

| | | |
|--|---|----------------------------------|
| for sinuses. Kerralite Cool Border As above with border. | 8 x 8cm 11 x 11cm 15 x 15cm | £2.02 £2.70 £4.29 |
| Medihoney (see under antimicrobials) | | |
| Flaminal Forte (note medihoney is cheaper than Flaminal) Alginate gel with 2 antimicrobial enzymes which do not damage healthy skin cells. Contains alginate. Debrides the wound and manages moisture balance. For wide range of wound types. Not for full thickness burns. Flaminal Hydro Contains lower concentration of alginate than Flaminal Forte | 15g 50g | £7.68 £25.42 |
| Hydrocoll Thin Thin transparent gelatin free hydrocolloid sheet with vapour permeable film backing. 7 days wear time. To promote epithelial growth. Conformable. | 7.5 x 7.5cm 10 x 10cm 15 x 15cm | 70p £1.16 £2.61 |
| Granuflex Border Hydrocolloid with adhesive foam border. Forms a gel on contact with exudate. Not vapour permeable. | 6 x 6cm 10 x 10cm 15 x 15cm 10 x 13cm sacral | £1.78 £3.37 £6.43 £3.97 |
| Debrisoft Debridement pad for removal of superficial slough and debris including biofilm. Do not apply with any emollients. Training required on use. | 10 x 10cm | £6.45 |
| OTHER ITEMS | | |
| Cellona Shoe (Activa Healthcare) For use when shoes do not fit due to bandaging or chronic oedema. | Petite -foot length up to 20cm Xsmall shoe size, 11.5-3, length 22.5cm Small, shoe size 3-6, length 24cm Medium, shoe size 6-7.5 length 26.5 Large, shoe size 7.5-9, length 28.5cm XL, shoe size 9-13, length 30cm | £15.75 |
| LimbO Waterproof Protector For showering and bathing over | Adult ½ leg Adult Elbow | £15.75 All sizes |

| | | |
|---|--|---|
| dressings. | | |
| Kerrapro Pressure reducing Pads Shaped pads 100% silicone, reusable, wash with soap and water, dry before re-application. Mimic fatty tissue and for redistributing pressure. Do not use on broken skin or over dressings. Best for category 1 pressure ulcers or bony prominences. | Sheet 10 x 10 x 0.3cm 10 x 10 x 1.2cm Strip 50 x 2.5 x 0.3cm 30 x 5 x 0.3cm Sacrum/ankle one size Heel one size | £4.24 £12.76 £5.31 £6.38 £16.85 £14.97 |