



York and Scarborough
Teaching Hospitals
NHS Foundation Trust

Circumcision for children

Information for parents, relatives and carers

Department of Urology

① For more information, please contact:

Paediatric Nurses

Telephone: 01904 726010

Ward 17

Telephone: 01904 726017

The York Hospital, Wigginton Road, York, YO31 8HE

Contents	Page
What is circumcision?	3
What happens before the operation?.....	4
What happens on the day of the operation?	5
What does the operation involve?	6
What happens after the operation?.....	6
Are there any risks in having a circumcision?	7
What are the alternatives to circumcision?	8
What issues should I look out for?	9
Tell us what you think of this leaflet	10
Teaching, training and research.....	10
Patient Advice and Liaison Service (PALS).....	10

What is circumcision?

Circumcision is the removal of the foreskin of the penis. The operation is usually performed for medical reasons, for example if the foreskin is too tight.

The operation is usually performed as a day patient when your child will usually spend a few hours at the hospital.

This leaflet explains a little about what will happen to your child before, during and after their operation. It tries to answer some of the questions you may have.

The doctors and nurses are there to help you and your child. They will always make time to listen to you and answer your questions. If you do not fully understand anything about your child's circumcision, please ask.

What happens before the operation?

You will have an appointment in a pre-assessment clinic. You will meet a nurse and a member of the play team. The nurse will ask medical questions and check the child's height and weight. The play team will discuss the anaesthetic to help you and the child understand it more. You will have the opportunity to ask any questions and be given some information to take home.

Please ensure you have some simple painkillers at home for after the operation, for example Calpol. If you are not sure what to buy, check with any pharmacy.

At some stage before or during the admission process, you will be asked to sign the second part of the consent form (FY03CON68-2 Circumcision (child) giving permission for the operation to take place, showing you understand what is to be done and confirming that you wish to proceed. The child can also sign the form if they wish. Make sure that you are given the opportunity to discuss any concerns and to ask any questions you may still have before signing the form.

What happens on the day of the operation?

You will have a letter that tells you either to go to Day Ward or Ward 17. These are both paediatric areas. The nurse will meet you and settle you into the ward. They will do some checks and give the child a name band. The doctor will explain the operation in more detail, discuss any worries you may have and ask you to sign a consent form. An anaesthetist will also see you to explain the child's anaesthetic in more detail. If the child has any medical problems or allergies, please tell the doctors.

The child can wear their own clothes to aid comfort. We recommend bringing loose comfortable clothing with no metalwork or zips. The ward can be quite warm.

After the operation the child will recover on the Day Ward or Ward 17. They need to stay a **minimum** of two hours on the ward but quite often they may need to stay longer. They can have a drink and then something to eat. We will ensure they have enough painkillers. We also like them to go to the toilet and pass urine before they go home.

If the child takes any regular medication, please bring this with you on the day of the operation

What does the operation involve?

Your child will have a general anaesthetic. This means they will be asleep during their circumcision. The anaesthetist will speak to you before their circumcision to discuss any concerns you may have and to check when they last had something to eat and drink.

The surgeon will also inject a local anaesthetic into the base of the penis to numb it. This will make it more comfortable for your child after the operation.

The surgeon trims the foreskin and inserts small dissolving stitches into the edges of the wound which may be supplemented with tissue glue.

What happens after the operation?

The child will benefit from extra rest for a couple of days. It is best to keep the child off school or nursery for two to three days. Please give painkillers according to the instructions on the packet to help ease pain. Sitting on toys or bicycles should be avoided until the wound has healed. Sporting activities such as PE and swimming can be restarted after two weeks.

Following the operation the wound should be kept clean and dry for 24 hours. The stitches are usually dissolvable and do not require removal. Bathing too early may cause the stitches to dissolve too quickly so we recommend quick washes for the first week after the operation.

You and the child may have an outpatient appointment several weeks after the surgery. If an appointment is required, a letter will be posted to you. Your nurse will give you discharge advice before you go home.

Are there any risks in having a circumcision?

As with any surgery, circumcision has some risks, which include:

Common (greater than one in 10)

- Swelling of the penis lasting several days

Occasional (between one in 10 and one in 50)

- Bleeding of the wound occasionally needing a further procedure
- Infection of the incision requiring further treatment and/or casualty visit
- Permanent altered or reduced sensation in the head of the penis
- Persistence of the absorbable stitches after three to four weeks, requiring removal

Rare (less than one in 50)

- Scar tenderness
- Failure to be completely satisfied with the cosmetic result
- Occasional need for removal of excessive skin at a later date
- Meatitis; inflammation of the urethral opening from the loss of protective foreskin which can lead to ulceration and meatal narrowing.

What are the alternatives to circumcision?

- Topical Steroids – The application of steroid cream to the foreskin can be beneficial.
- Stretching Techniques – Can be useful for the non-retractable foreskin when scarring is not present.
- Conservative Surgery - A number of plastic surgical techniques are available for the tight foreskin of which the commonest is prepuceplasty. This procedure is less invasive and has fewer complications than circumcision and allows the foreskin to be retained.

What issues should I look out for?

- The child is not drinking well at home.
- The child is in a lot of pain and painkillers do not seem to help.
- The operation site seems inflamed, red, or the skin seems hotter than the surrounding skin.
- There is any oozing from the operation site.

If you have these or other concerns after the operation please use the following numbers:

Monday to Friday, day time: 01904 726010.

Please ask to speak to a paediatric nurse.

At other times, or if no paediatric nurse is available at the above number: 01904 722017.

Please note the nurses may give you advice, or they may direct you to contact your GP.

Tell us what you think of this leaflet

Meeting the needs and preferences of patients and carers is at the centre of everything we do. We hope that you found this leaflet useful and informative.

If you would like to comment on it, please contact Mr Mustafa Hilmy, Consultant Urological Surgeon, Department of Urology, The York Hospital, Wigginton Road, York, YO31 8HE, telephone 01904 725846 or email Lisa.Browne@york.nhs.uk.

Teaching, training and research

Our Trust is committed to teaching, training and research to support the development of health and healthcare in our community. Healthcare students may observe consultations for this purpose. You can opt out if you do not want students to observe. We may also ask you if you would like to be involved in our research.

Patient Advice and Liaison Service (PALS)

PALS offers impartial advice and assistance to patients, their relatives, friends and carers. We can listen to feedback (positive or negative), answer questions and help resolve any concerns about Trust services.

PALS can be contacted on 01904 726262, or email pals@york.nhs.uk.

An answer phone is available out of hours.

Leaflets in alternative languages or formats

Please telephone or email if you require this information in a different language or format, for example Braille, large print or audio.

如果你要求本資 不同的 或 式提供 , 電
或發電

Jeżeli niniejsze informacje potrzebne są w innym języku lub formacie, należy zadzwonić lub wysłać wiadomość e-mail

Bu bilgileri değişik bir lisanda ya da formatta istiyorsanız lütfen telefon ediniz ya da e-posta gönderiniz

Telephone: 01904 725566

Email: access@york.nhs.uk

Owner	Mr Hilmy, Consultant Urologist
Date first issued	November 2002
Review Date	April 2024
Version	8 (issued April 2021)
Approved by	Urology clinical governance meeting
Linked to consent form	FY03CON68-2 Circumcision (child) v4
Document Reference	PIL 141 v8
© 2021 York and Scarborough Teaching Hospitals NHS Foundation Trust. All Rights reserved.	